

Patient Assessment Scenario "I" Testing Form



GROUP:		LEARNER:		☐ PRACTICE ☐ TEST ☐ REMEDIAL	
INSTRUCTOR:		FEEDBACK PROVIDER:		ASSESSOR:	

 CRITICAL TREATMENT DECISIONS (Check Yes or No): The Learner must:	Y	N
1. Performed the assessment and management phases properly and in correct sequence.	☐	☐
2. Recognized that respiratory distress is related shock and pneumothorax, not an obstructed airway.	☐	☐
3. Performed tube thoracostomy.	☐	☐
4. Reassessed the xABCDEs and established a differential diagnosis for continued shock.	☐	☐
5. Identified cardiac tamponade as the likely cause of continued shock and obtained a surgical consult for definitive treatment.	☐	☐
6. Performed volume resuscitation as a temporizing measure while awaiting surgical intervention.	☐	☐

Potentials for Adverse Outcome: If the Learner's performance is borderline and the Learner fails to recognize or perform one or more of the items below, the Instructor may determine that the Learner's overall performance is unsuccessful. OBSERVED LIST OF MANEUVERS (DECISIONS) (Check Yes or No):	Y	N
1. Avoided rapid sequence intubation which would result in cardiac arrest.	☐	☐
2. Recognized distended neck veins.	☐	☐
3. Did not order extensive imaging.	☐	☐
4. Did not delay thoracostomy for imaging.	☐	☐
5. Avoided pericardiocentesis.	☐	☐
6. Followed needle or finger thoracostomy with tube thoracostomy.	☐	☐
7. Did not delay surgical consultation for imaging results.	☐	☐
8. Considered tetanus immunization and antibiotics.	☐	☐

RATING:	☐ SUCCESSFUL	☐ INCOMPLETE/REMEDIAL	☐ INCOMPLETE/REPEAT COURSE
REMEDIAL RATING:	☐ SUCCESSFUL	☐ INCOMPLETE/REPEAT COURSE	
INSTRUCTOR POTENTIAL?	☐ YES	☐ NO	

Comments:

